



— NOW APPROVED —

CODING AND REIMBURSEMENT GUIDE

ICD-10-CM Diagnosis, HCPCS, CPT®, and NDC Billing Information
for ANDEMBRY® (garadacimab-gxii)

CODING INFORMATION FOR ANDEMBRY® (garadacimab-gxii)

Disclaimers

- All codes provided are for informational purposes only and are subject to change without notice
- The treating physician is solely responsible for diagnosis coding and determination of the appropriate ICD-10-CM codes that describe the patient’s condition and are supported by medical records
- Please contact third-party payers directly for specific information on their coding, coverage, payment policies, and fee schedules

For questions regarding reimbursement support, please call ANDEMBRY ConnectSM at 1-844-423-4273

ICD-10-CM Diagnosis Code for Persons With Hereditary Angioedema¹

ICD-10-CM Code	Description
D84.1	Defects in the complement system C1 esterase inhibitor (C1-INH) deficiency

HCPCS Codes²

The following drug codes have been assigned by the Center for Medicare and Medicaid Services (CMS) National Level II HCPCS coding system and can be used for ANDEMBRY. The provider is advised to verify the correct codes directly with the third-party payer.

HCPCS Code	Description
J3490	Unclassified drugs
J3590	Unclassified biologics

Note: These HCPCS codes are temporary and will be replaced by a permanent J-code once it is available.

HCPCS, Healthcare Common Procedure Coding System; ICD-10-CM, International Classification of Diseases, Tenth Revision, Clinical Modification.

Please see Important Safety Information on page 4 and accompanying full Prescribing Information.

CPT® Administration Codes: Physician Office³

ANDEMBRY is a subcutaneous injection that may be self-administered or administered by a caregiver via a prefilled autoinjector subcutaneously into the thigh or abdomen, or to the upper arm if a caregiver administers the injection.

For Medicare beneficiaries, the following codes may not be payable. Medicare considers ANDEMBRY self-administered and therefore not covered as a Part B service. However, other payers may allow administration in a physician office or other site of service. The healthcare professional may not bill for the administration of ANDEMBRY if provided through the QuickStart Program.

Please verify current coding requirements directly with the payer being billed.

CPT® Code	Description
96372	Therapeutic, prophylactic, or diagnostic injection (specify substance or drug); subcutaneous or intramuscular

Do not report 96372 for injections given without direct physician or other qualified health professional supervision. To report, use 99211.

Note from CPT National Correct Coding Initiative (NCCI) Policy Manual for Medicare Services: CPT code 96372 is reportable by physicians (MD, DO, or other practitioners who bill Medicare contractors) for services performed in physicians’ offices. These drug administration services should not be reported by physicians for services provided in a facility setting such as a hospital outpatient, ambulatory surgery center, or emergency department.⁴

CPT® Code	Description
99211	Office or other outpatient visit for the evaluation and management of an established patient that may not require the presence of a physician or other qualified healthcare professional. Usually, the presenting problem(s) are minimal. Typically, 5 minutes are spent performing or supervising these services.

Note from CPT NCCI Policy Manual for Medicare Services: Drug administration CPT code 96372 has been valued to include the work and practice expenses of CPT code 99211. Therefore, CPT code 99211 is not reportable with CPT code 96372. Other non-facility–based evaluation and management codes may be separately reportable with modifier 25 if the physician provides a significant and separately identifiable E&M service.⁴

ANDEMBRY National Drug Codes (NDCs)

NDC (Device)	NDC (Carton)	Strength	Presentation
63833-925-20	63833-925-01	200 mg/1.2 mL	Prefilled autoinjector

CPT, current procedural terminology.

CPT © 2025 American Medical Association. All rights reserved.



ANDEMBRY® (garadacimab-gxii) CMS-1500 CLAIM FORM

HEALTH INSURANCE CLAIM FORM

APPROVED BY NATIONAL UNIFORM CLAIM COMMITTEE (NUCC) 02/12

PICA										PICA									
1. MEDICARE (Medicare#) MEDICAID (Medicaid#) TRICARE (ID#/DoD#) CHAMPVA (Member ID#) GROUP HEALTH PLAN (ID#) FECA BLK LUNG (ID#) OTHER (ID#)										1a. INSURED'S I.D. NUMBER (For Program in Item 1)									
2. PATIENT'S NAME (Last Name, First Name, Middle Initial)										3. PATIENT'S BIRTH DATE MM DD YY SEX M F									
5. PATIENT'S ADDRESS (No., Street)										6. PATIENT RELATIONSHIP TO INSURED Self Spouse Child Other									
CITY STATE										7. INSURED'S ADDRESS (No., Street)									
ZIP CODE TELEPHONE (Include Area Code)										CITY STATE									
9. OTHER INSURED'S NAME (Last Name, First Name, Middle Initial)										10. IS PATIENT'S CONDITION RELATED TO:									
a. OTHER INSURED'S POLICY OR GROUP NUMBER										a. EMPLOYMENT? (Current or Previous) YES NO									
b. RESERVED FOR NUCC USE										b. AUTO ACCIDENT? PLACE (State) YES NO									
c. RESERVED FOR NUCC USE										c. OTHER ACCIDENT? YES NO									
d. INSURANCE PLAN NAME OR PROGRAM NAME										10d. CLAIM CODES (Designated by NUCC)									
12. PATIENT'S OR AUTHORIZED PERSON'S SIGNATURE I authorize the release of any medical or other information necessary to process this claim. I also request payment of government benefits either to myself or to the party who accepts assignment below.										13. INSURED'S OR AUTHORIZED PERSON'S SIGNATURE I authorize payment of medical benefits to the undersigned physician or supplier for services described below.									
SIGNED DATE										SIGNED									
14. DATE OF CURRENT ILLNESS, INJURY, OR PREGNANCY (LMP) MM DD YY QUAL										15. OTHER DATE MM DD YY QUAL									
17. NAME OF REFERRING PROVIDER OR OTHER SOURCE										17a. 17b. NPI									
19. ADDITIONAL CLAIM INFORMATION (Designated by NUCC)										16. DATES PATIENT UNABLE TO WORK IN CURRENT OCCUPATION FROM TO MM DD YY MM DD YY									
21. DIAGNOSIS OR NATURE OF ILLNESS OR INJURY Relate A-L to service line below (24E)										18. HOSPITALIZATION DATES RELATED TO CURRENT SERVICES FROM TO MM DD YY MM DD YY									
A. B. C. D. ICD Ind.										20. OUTSIDE LAB? YES NO \$ CHARGES									
E. F. G. H.										22. RESUBMISSION CODE ORIGINAL REF. NO.									
I. J. K. L.										23. PRIOR AUTHORIZATION NUMBER									
24. A. DATE(S) OF SERVICE From To MM DD YY MM DD YY B. PLACE OF SERVICE C. EMG D. PROCEDURES, SERVICES, OR SUPPLIES (Explain Unusual Circumstances) E. DIAGNOSIS F. \$ CHARGES G. DAYS OR UNITS H. EPOT Family Plan I. ID. QUAL J. RENDERING PROVIDER ID. #																			
1										D E F									
2																			
3																			
4																			
5																			
6																			
25. FEDERAL TAX I.D. NUMBER SSN EIN										26. PATIENT'S ACCOUNT NO.									
31. SIGNATURE OF PHYSICIAN OR SUPPLIER INCLUDING DEGREES OR CREDENTIALS (I certify that the statements on the reverse apply to this bill and are made a part thereof.)										27. ACCEPT ASSIGNMENT? (For govt. claims, see back) YES NO									
32. SERVICE FACILITY LOCATION INFORMATION										28. TOTAL CHARGE \$ 29. AMOUNT PAID \$ 30. Ptsd. for NUCC Use									
33. BILLING PROVIDER INFO & PH#																			
SIGNED DATE										a. NPI b.									

NUCC Instruction Manual available at: www.nucc.org

PLEASE PRINT OR TYPE

APPROVED OMB-0938-1197 FORM 1500 (02-12)

A Field 19: Additional Claim Information²

When an unclassified drug HCPCS code (J3490 or J3590) is used, the provider must include additional information about ANDEMBRY in a free-text field, such as Field 19 or its electronic equivalent. Payers typically require the name of the drug, its NDC, the route of administration, and the amount used for the treatment. For example:

- ANDEMBRY NDC (Device) 63833-925-20; Prefilled autoinjector; 200 mg/1.2 mL
- ANDEMBRY NDC (Carton) 63833-925-01; Prefilled autoinjector; 200 mg/1.2 mL

B Field 21: Diagnosis or Nature of Illness or Injury

Enter the ICD-10-CM diagnosis code(s) that best describes the reason the patient is receiving care.

C **Field 24A (Shaded Area): NDC**

For Medicaid claims and for Medicare claims that will cross over to Medicaid as the secondary payer, NDC information in a specific format is required in the shaded area above the line on which ANDEMBRY is reported in Field 24A. Various state Medicaid plans and Medicare have different reporting formats for this information.

Other payers may require different information. Check with the payer for specific requirements.

D Field 24D: CPT/HCPCS^{2,3}

HCPCS coding options for ANDEMBRY are:

- J3490 Unclassified drugs
- J3590 Unclassified biologics

For Medicare beneficiaries, the following code may not be payable:

- 96372: Therapeutic, prophylactic, or diagnostic injection (specify substance or drug); subcutaneous or intramuscular

The code selected is based on the documented service provided to the patient.

Medicare considers ANDEMBRY self-administered and therefore not covered as a Part B service. However, other payers may allow administration in a physician office or other site of service. Please verify current coding requirements directly with the payer being billed.

E Field 24E: Diagnosis Pointer

Enter the line number(s) from Field 21 that best describes the medical necessity for the service.

F	Field 24G: Days or Units
---	--------------------------

Requirements for Field 24G vary by payer. For example, some payers rely on the information in Field 19 for these data and request a placeholder of "1" in Field 24G. Other payers may require 1 unit per unit of drug used in the care of the patient.

Important Safety Information

ANDEMBRY® (garadacimab-gxii) injection, for subcutaneous use, is an activated Factor XII (FXIIa) inhibitor (monoclonal antibody) indicated for prophylaxis to prevent attacks of hereditary angioedema (HAE) in adult and pediatric patients aged 12 years and older.

The most common adverse reactions in the pivotal trial (incidence ≥7%) were nasopharyngitis and abdominal pain. In the pivotal trial and the open-label extension study, injection-site reactions (eg, injection-site bruising, injection-site erythema, injection-site hematoma, injection-site pruritus, injection-site urticaria) were reported in 23 (14%) patients.

No dedicated drug interaction studies have been conducted, nor is there data concerning the use of ANDEMBRY in women who are pregnant or breastfeeding. The safety and efficacy of ANDEMBRY in patients under 12 years of age have not been established.

Drug Interference with Laboratory Test: ANDEMBRY can prolong activated partial thromboplastin time (aPTT) due to an interaction of garadacimab-gxii with the aPTT assay.

Please see full Prescribing Information for ANDEMBRY.

To report SUSPECTED ADVERSE REACTIONS, contact the CSL Behring Pharmacovigilance Department at 1-866-915-6958 or FDA at 1-800-FDA-1088 or www.fda.gov/medwatch.

Please see accompanying full Prescribing Information.



ANDEMBRY Connect provides ongoing, one-on-one support throughout the treatment journey

- Personalized support for each patient
- Dedicated case manager for each office
- Injection training and product support available through our dedicated nursing team
- Financial assistance*
- Benefits verification and prior authorization support (if applicable)

QuickStart Program

- ANDEMBRY QuickStart Program for commercially and government-insured patients

*Subject to terms and conditions of the copay assistance program. To read the full terms and conditions, visit <http://cslbehring.com/patients/support-and-assistance>. CSL Behring reserves the right to rescind, revoke, or amend the program at any time without notice.



Call **[1-844-423-4273]**
for ANDEMBRY Connect from **[8 AM to 8 PM ET]** Monday–Friday

References: 1. US Department of Health and Human Services, Centers for Medicare & Medicaid Services. International Classification of Diseases, Tenth Revision, Clinical Modification (ICD-10-CM). Accessed April 17, 2025. <https://www.cms.gov/medicare/coding-billing/icd-10-codes> 2. US Department of Health and Human Services, Centers for Medicare and Medicaid Services. HCPCS release and code sets. Accessed April 17, 2025. <https://www.cms.gov/medicare/coding-billing/healthcare-common-procedure-system/quarterly-update> 3. American Academy of Professional Coders. Accessed April 17, 2025. <https://coder.aapc.com/cpt-codes> 4. Medicaid NCCI 2023 Coding Policy Manual Chapter 11. Accessed April 17, 2025. <https://www.cms.gov/files/document/medicaid-ncci-policy-manual-2024-chapter-11.pdf>

CSL Behring

ANDEMBRY is manufactured by CSL Behring GmbH and distributed by CSL Behring LLC.
ANDEMBRY® is a registered trademark of CSL Behring LLC.
ANDEMBRY ConnectSM is a service mark of CSL Behring LLC.
All other trademarks are property of their respective owners.
©2025 CSL Behring LLC. 1020 First Avenue, PO Box 61501, King of Prussia, PA 19406-0901 USA
www.CSLBehring.com USA-AND-0028-JUN25



HIGHLIGHTS OF PRESCRIBING INFORMATION
These highlights do not include all the information needed to use ANDEMBRY safely and effectively. See full prescribing information for ANDEMBRY.

ANDEMBRY® (garadacimab-gxii) injection, for subcutaneous use
Initial U.S. Approval: 2025

INDICATIONS AND USAGE
ANDEMBRY is an activated Factor XII (FXIIa) inhibitor (monoclonal antibody) indicated for prophylaxis to prevent attacks of hereditary angioedema (HAE) in adult and pediatric patients aged 12 years and older. (1)

DOSAGE AND ADMINISTRATION

- Recommended Dosage: Initial loading dose of 400 mg (two 200 mg injections) administered subcutaneously followed by maintenance dosage of 200 mg once monthly. (2.1)
- Subcutaneous use only. (2.2)
- Patients may self-administer. See full prescribing information for preparation and administration instructions. (2.2)

DOSAGE FORMS AND STRENGTHS
Injection:

- 200 mg/1.2 mL solution in single-dose prefilled autoinjector (3)
- 200 mg/1.2 mL (167 mg/mL) solution in single-dose prefilled syringe with needle safety device (3)

CONTRAINDICATIONS
None. (4)

WARNINGS AND PRECAUTIONS
None. (5)

ADVERSE REACTIONS
Most common adverse reactions (incidence $\geq 7\%$) are nasopharyngitis and abdominal pain (6.1)

To report SUSPECTED ADVERSE REACTIONS, contact CSL Behring at 1-866-915-6958 or FDA at 1-800-FDA-1088 or www.fda.gov/medwatch.

DRUG INTERACTIONS
Drug Interference with Laboratory Test: ANDEMBRY can prolong activated partial thromboplastin time (aPTT) due to an interaction of garadacimab-gxii with the aPTT assay. (7.1)

See 17 for PATIENT COUNSELING INFORMATION and FDA-approved patient labeling.

Revised: June 2025

FULL PRESCRIBING INFORMATION: CONTENTS*

- 1 INDICATIONS AND USAGE**
- 2 DOSAGE AND ADMINISTRATION**
 - 2.1 Recommended Dosage
 - 2.2 Preparation and Administration Instructions for ANDEMBRY Prefilled Autoinjector and Prefilled Syringe with Needle Safety Device
- 3 DOSAGE FORMS AND STRENGTHS**
- 4 CONTRAINDICATIONS**
- 5 WARNINGS AND PRECAUTIONS**
- 6 ADVERSE REACTIONS**
 - 6.1 Clinical Trials Experience
- 7 DRUG INTERACTIONS**
 - 7.1 Drug Interference with Laboratory Test
- 8 USE IN SPECIFIC POPULATIONS**
 - 8.1 Pregnancy
 - 8.2 Lactation
 - 8.4 Pediatric Use
 - 8.5 Geriatric Use

- 11 DESCRIPTION**
- 12 CLINICAL PHARMACOLOGY**
 - 12.1 Mechanism of Action
 - 12.2 Pharmacodynamics
 - 12.3 Pharmacokinetics
 - 12.6 Immunogenicity
- 13 NONCLINICAL TOXICOLOGY**
 - 13.1 Carcinogenesis, Mutagenesis, Impairment of Fertility
- 14 CLINICAL STUDIES**
- 16 HOW SUPPLIED/STORAGE AND HANDLING**
- 17 PATIENT COUNSELING INFORMATION**

* Sections or subsections omitted from the full prescribing information are not listed

CSL Behring

ANDEMBRY®

(garadacimab-gxii) injection, for subcutaneous use

FULL PRESCRIBING INFORMATION

1 INDICATIONS AND USAGE

ANDEMBRY is indicated for prophylaxis to prevent attacks of hereditary angioedema (HAE) in adult and pediatric patients aged 12 years and older.

2 DOSAGE AND ADMINISTRATION

2.1 Recommended Dosage

The recommended dosage of ANDEMBRY is an initial loading dose of 400 mg (two injections of 200 mg) administered subcutaneously on the first day of treatment followed by a maintenance dosage of 200 mg administered subcutaneously every month.

Missed Dose(s)

If a dose of ANDEMBRY is missed, administer the dose as soon as possible.

2.2 Preparation and Administration Instructions for ANDEMBRY Prefilled Autoinjector and Prefilled Syringe with Needle Safety Device

- For subcutaneous use only.
- ANDEMBRY is intended for self-administration or administration by a caregiver. Prior to treatment initiation, train patients/caregivers on proper preparation and subcutaneous (SC) administration technique of ANDEMBRY [see *Instructions for Use*].
- Prior to administration, remove ANDEMBRY from the refrigerator and allow to sit for 30 minutes at room temperature before use.
- Inspect ANDEMBRY visually for particulate matter and discoloration prior to administration, whenever solution and container permit. ANDEMBRY is a slightly opalescent to clear, brownish-yellow to yellow solution.
- Administer ANDEMBRY subcutaneously into the thigh or abdomen ensuring to stay 1 inch (2 cm) away from the navel. The upper arm can also be used if a caregiver administers the subcutaneous injection.
- Discard the used ANDEMBRY into a sharps disposal container (closed puncture-resistant container).
- For detailed instructions on the preparation and administration of ANDEMBRY [see *Instructions for Use*].

3 DOSAGE FORMS AND STRENGTHS

Injection: a sterile, preservative-free, slightly opalescent to clear, brownish-yellow to yellow solution available in the following:

- 200 mg/1.2 mL of garadacimab solution in a single-dose prefilled autoinjector
- 200 mg/1.2 mL (167 mg/mL) of garadacimab solution in a single-dose prefilled syringe with needle safety device

4 CONTRAINDICATIONS

None.

5 WARNINGS AND PRECAUTIONS

None.

6 ADVERSE REACTIONS

6.1 Clinical Trials Experience

Because clinical trials are conducted under widely varying conditions, adverse reaction rates observed in the clinical trials of a drug cannot be directly compared to rates in the clinical trials of another drug and may not reflect the rates observed in practice.

The safety of ANDEMBRY reflects the exposure in a total of 164 adult and pediatric patients aged 12 years and older with hereditary angioedema (HAE) from a placebo-controlled study, VANGUARD [see *Clinical Studies* (14)], and an open-label clinical study. Among the 164 patients who received at least one dose of ANDEMBRY 200 mg subcutaneously, 153 (93%) patients were exposed for at least one year. The median duration of ANDEMBRY treatment was 2.6 years.

The safety data below is based on the 6-month placebo-controlled study (VANGUARD), in which ANDEMBRY 400 mg was administered subcutaneously as a loading dose followed by 200 mg (N=39) every month in patients with HAE. Demographics of the patients in this study are summarized in Clinical Studies [see *Clinical Studies* (14)]. The safety of ANDEMBRY was similar across all subgroups of patients, including analysis by age, sex and geographic region.

Table 1 provides the most common adverse reactions with ANDEMBRY with incidence $\geq 7\%$ and more common than placebo.

Table 1 Adverse Reactions with ANDEMBRY with Incidence $\geq 7\%$ and More Common than Placebo in Patients with HAE (VANGUARD)

Adverse Reactions	ANDEMBRY (N=39)	Placebo (N=25)
	n (%)	n (%)
Nasopharyngitis*	8 (21)	3 (12)
Abdominal Pain**	3 (8)	0

*Consists of nasopharyngitis, rhinitis, and upper respiratory infections

**Consists of abdominal pain and abdominal pain lower

Specific Adverse Reaction(s):

Injection Site Reactions

In VANGUARD and an open-label clinical study, which included 57 patients who rolled over from VANGUARD, 164 patients with HAE received ANDEMBRY 200 mg subcutaneously every month. Injection site reactions (e.g., injection site bruising, injection site erythema, injection site hematoma, injection site pruritus, injection site urticaria) were reported in 23 (14%) patients.

Laboratory Abnormalities:

Prolonged Coagulation Tests (aPTT and PT)

In the VANGUARD trial, the incidence of prolonged activated partial thromboplastin time (aPTT), defined as $>1.4 \times \text{ULN}$, was 3 (8%) patients in the ANDEMBRY group compared to 0 patients in the placebo group. Additionally, the incidence of prolonged prothrombin time (PT) or international normalized ratio (INR), defined as $>1.3 \times \text{ULN}$, was 6 (15%) patients in the ANDEMBRY group compared to 1 (4%) patient in the placebo group. None of the increases in aPTT, PT and INR were associated with bleeding events [see *Drug Interactions* (7.1)].

7 DRUG INTERACTIONS

7.1 Drug Interference with Laboratory Test

Coagulation Tests

ANDEMBRY can prolong activated partial thromboplastin time (aPTT) due to an interaction of garadacimab-gxii with the aPTT assay. The reagents used in the aPTT laboratory test initiate intrinsic coagulation through the activation of FXII in the contact system, therefore inhibition of plasma FXIIa by ANDEMBRY can prolong aPTT in this assay [see *Adverse Reactions* (6.1)].

8 USE IN SPECIFIC POPULATIONS

8.1 Pregnancy

Risk Summary

There are no available data on ANDEMBRY use in pregnant women to evaluate for a drug-associated risk of major birth defects, miscarriage or other adverse maternal or fetal outcomes. Monoclonal antibodies such as garadacimab-gxii are transported across the placenta during the third trimester of pregnancy; therefore, potential effects on a fetus are likely to be greater during the third trimester of pregnancy. In an embryo-fetal development study in pregnant rabbits, garadacimab-gxii produced no evidence of fetal harm at doses up to approximately 100 times the exposure achieved at the maximum recommended human dose (MRHD) of 200 mg once monthly. In a pre- and post-natal development study in rabbits, garadacimab-gxii had no effects on survival, growth, or development of F₁ offspring at doses resulting in approximately 76 times the exposure achieved at the MRHD (see *Data*).

The estimated background risk of major birth defects and miscarriage for the indicated population(s) is unknown. All pregnancies have a background risk of birth defect, loss, or other adverse outcomes. In the U.S. general population, the estimated background risk of major birth defects and miscarriage in clinically recognized pregnancies is 2% to 4% and 15% to 20%, respectively.

Data

Animal Data

In an embryo-fetal development study in pregnant rabbits dosed by the intravenous route every 3 days during the period of organogenesis from gestation days 6 to 18, garadacimab-gxii produced no evidence of fetal harm or maternal toxicity at doses resulting in approximately 100 times the exposure achieved at the MRHD (on an AUC basis with maternal intravenous doses up to 100 mg/kg/dose).

In a pre- and post-natal development study in rabbits dosed by the intravenous or subcutaneous route once every 5 days from the end of implantation (Gestation Day 7) to postpartum day 38 (end of the lactation period), garadacimab-gxii had no effects on survival, growth, or development of F₁ offspring at doses resulting in approximately 76 times the exposure achieved at the MRHD (on an AUC basis with a maternal intravenous dose of 100 mg/kg/dose). There was no evidence of maternal toxicity.

Garadacimab-gxii crossed the placenta in rabbits. With a maternal dose of 100 mg/kg garadacimab-gxii on gestation day 29, fetal plasma concentrations were 40.8% of maternal concentrations following subcutaneous administration and approximately equivalent following intravenous administration.

8.2 Lactation

Risk Summary

There are no data on the presence of garadacimab-gxii or its metabolite in either human or animal milk, the effects on the breastfed infant, or the effects on milk production. Maternal IgG is known to be present in human milk. The effects of local gastrointestinal exposure and limited systemic exposure in the breastfed infant to garadacimab-gxii are unknown. The developmental and health benefits of breastfeeding should be considered along with the mother's clinical need for ANDEMBRY and any potential adverse effects on the breastfed infant from ANDEMBRY or the underlying maternal condition.

8.4 Pediatric Use

The safety and effectiveness of ANDEMBRY for prophylaxis to prevent attacks of HAE have been established in pediatric patients aged 12 years and older. Use of ANDEMBRY for this indication is supported by evidence from a total of 6 pediatric patients aged 12 years and older enrolled in VANGUARD who received treatment with ANDEMBRY 400 mg loading dose administered subcutaneously followed by a maintenance dosage of 200 mg subcutaneously every month (n=4) or placebo (n=2). Results of the subgroup analysis for pediatric patients aged 12 years and older were consistent with study results for adult patients [see *Adverse Reactions* (6.1), *Clinical Pharmacology* (12.3), and *Clinical Studies* (14)].

The safety and effectiveness of ANDEMBRY have not been established in pediatric patients younger than 12 years of age.

8.5 Geriatric Use

Of the total number of ANDEMBRY-treated patients in the clinical study for prophylaxis to prevent attacks of HAE, 6 (9%) were 65 years of age and older [see *Clinical Studies* (14)]. No overall differences in safety or effectiveness of ANDEMBRY have been observed between patients 65 years of age and older and younger adult patients [see *Adverse Reactions* (6.1), *Clinical Pharmacology* (12.3), and *Clinical Studies* (14)].

11 DESCRIPTION

Garadacimab-gxii, an activated Factor XII (FXIIa) inhibitor, is a recombinant, fully human, monoclonal antibody (IgG4/ λ -light chain) produced in Chinese Hamster Ovary (CHO) cells. Based on the amino acid sequence, the molecular weight of garadacimab-gxii is approximately 148 kDa.

ANDEMBRY (garadacimab-gxii) injection is a sterile, preservative-free, slightly opalescent to clear, brownish-yellow to yellow solution for subcutaneous use.

Each 1.2 mL prefilled autoinjector or prefilled syringe with needle safety device of ANDEMBRY contains 200 mg of garadacimab-gxii, arginine monohydrochloride (37.9 mg),

histidine (3.7 mg), polysorbate 80 (0.24 mg), proline (19.3 mg), and water for injection, USP. The pH is 6.1.

12 CLINICAL PHARMACOLOGY

12.1 Mechanism of Action

Garadacimab-gxii is an inhibitor of activated FXII that binds to the catalytic domain of activated Factor XII (FXIIa and βFXIIa) and inhibits its catalytic activity. FXII is the first factor activated in the contact activation pathway and initiates the inflammatory bradykinin-producing kallikrein-kinin system. The inhibition of FXIIa decreases the activation of prekallikrein to kallikrein and the generation of bradykinin, which is associated with inflammation and swelling in HAE attacks, thus reducing the cascade of events leading to an HAE attack.

12.2 Pharmacodynamics

Garadacimab-gxii concentration-dependent inhibition of FXIIa-mediated kallikrein activity was observed in patients with HAE. Inhibition of FXIIa-mediated kallikrein activity was observed after the first dose.

12.3 Pharmacokinetics

The pharmacokinetics of garadacimab-gxii are provided in Table 2 following subcutaneous administration of a single 200 mg dose to healthy volunteers and are presented as mean (SD), unless otherwise specified.

Table 2 Garadacimab-gxii Pharmacokinetics Following Subcutaneous Administration*

Parameter	Garadacimab-gxii
Exposure	
C _{max}	18.4 (7.23) mcg/mL
AUC _{0-inf}	11470 (4103) mcg·h/mL
C _{trough, Steady State} ^{a,b}	8.09 (4.28) to 8.69 (3.93) mcg /mL
Absorption	
Absolute Bioavailability	39 (34, 43)%
T _{max}	6 (2, 15) days
Distribution	
Apparent Volume of Distribution	11.8 (5.17) Liters (L)
Elimination	
Half-Life ^b	17.4 (3.14) days
Apparent Clearance	0.02 (0.008) L/h
Metabolism	
Primary Pathway	Expected to be metabolized into small peptides by catabolic pathways

Abbreviations: AUC = area under the plasma concentration-time curve; CI = confidence interval; C_{max} = maximum plasma concentration; C_{trough, steady state} = trough plasma concentration at steady state; T_{max} = time to peak concentration
T_{max} is reported as median (range) and absolute bioavailability is reported as estimate (95% CI)
* The PK of garadacimab-gxii is similar between healthy volunteers and subjects with HAE
^a HAE patients enrolled in the VANGUARD trial
^b Garadacimab-gxii exposure (based on AUC_{0-inf} and C_{max}) following the initial subcutaneous administration of loading dose of 400 mg (2 doses of 200 mg) is 133% of steady-state exposures following 200 mg subcutaneous once monthly.

Specific Populations

No clinically significant differences in the pharmacokinetics of garadacimab-gxii were observed based on age (age range: 12 to 73 years), gender, race, body weight (range: 43 to 153 kg), and mild to moderate renal impairment (eGFR 30 to <90 mL/min/1.73 m²). The effect of severe renal impairment (eGFR <30 mL/min/1.73 m²) on garadacimab-gxii pharmacokinetics is unknown.

Pediatric Patients

No clinically significant difference in garadacimab-gxii C_{max} and AUC_{tau} was observed between pediatric patients (age range: 12 to 17 years) and adults who received the recommended dosage of garadacimab-gxii (200 mg of garadacimab-gxii once monthly).

Drug Interactions

No dedicated drug interaction studies have been conducted in humans.

12.6 Immunogenicity

The observed incidence of anti-drug antibodies (ADA) is highly dependent on the sensitivity and specificity of the assay. Differences in assay methods preclude meaningful comparisons of the incidence of anti-drug antibodies in the studies described below with the incidence of anti-drug antibodies in other studies, including those of garadacimab-gxii or other garadacimab products.
In VANGUARD, 2.6% (1/39) of patients who received garadacimab-gxii tested positive for anti- garadacimab-antibodies during the 6-month treatment period.
The development of ADA against garadacimab-gxii did not affect pharmacokinetics (PK), safety, or clinical response.

13 NONCLINICAL TOXICOLOGY

13.1 Carcinogenesis, Mutagenesis, Impairment of Fertility

Animal studies have not been conducted to evaluate the carcinogenic or mutagenic potential of garadacimab-gxii.
Fertility and reproductive performance were unaffected in sexually mature male and female rabbits that received garadacimab-gxii at intravenous doses up to 100 mg/kg once every three days (resulting in exposures in male and female rabbits up to approximately 100 and 80 times the MRHD on an AUC basis).

14 CLINICAL STUDIES

The efficacy of ANDEMBRY for prophylaxis to prevent hereditary angioedema (HAE) attacks was evaluated in a multicenter, randomized, double-blind, placebo-controlled, parallel-group study (VANGUARD [NCT04656418]) of 6 months duration.

VANGUARD enrolled a total of 64 adult and pediatric patients 12 years of age and older with Type I or Type II HAE who experienced at least 2 investigator-confirmed HAE attacks over a 2-month period prior to treatment with ANDEMBRY or placebo. Patients were randomized in a 3:2 ratio (ANDEMBRY:placebo) to receive an initial loading dose of ANDEMBRY 400 mg administered subcutaneously followed by a maintenance dose of 200 mg every month, or matched placebo. Patients were required to discontinue other prophylactic HAE medications prior to entering the study. All patients were allowed to use on-demand medications for the treatment of HAE attacks during the study.
The demographics and baseline characteristics of patients in VANGUARD are provided in Table 3.

Table 3 Demographics and Baseline Characteristics of Patients in the VANGUARD Trial

	VANGUARD (N=64)
Female, n (%)	38 (59)
Race, n (%)	
Asian	6 (9)
Black or African American	1 (2)
Native Hawaiian or Pacific Islander	1 (2)
White	55 (86)
Other	1 (2)
Hispanic or Latino, n (%)	3 (5)
Mean age, years (SD)	41 (16)
≤ 17 years, n (%)	6 (9)
> 17 years, n (%)	58 (91)
C1-INH HAE Type 1, n (%)	56 (88)
Prior HAE prophylaxis [‡] , n (%)	21 (33)
Baseline HAE attack rate*, n (%)	
1 to < 3 attacks	26 (41)
≥ 3 attacks	38 (59)

* Within 3 months before screening
[‡] During 2-month period prior to treatment

The primary endpoint for VANGUARD was the monthly HAE attack rate at 6 months (number of investigator-confirmed HAE attacks per month). As shown in Table 4, the least squares mean for the monthly HAE attack rate was lower with ANDEMBRY compared to placebo.

Table 4 Rate of Monthly HAE Attack at 6 Months in VANGUARD

Rate of Monthly HAE Attack	ANDEMBRY (N=39)	Placebo (N=25)
LS mean (95% CI)	0.22 (0.11, 0.47)	2.07 (1.49, 2.87)
Percent reduction relative to placebo (95% CI)	89.2 (75.6, 95.2)	
P-value ^a	< 0.001	

Abbreviations: CI=confidence interval; HAE=hereditary angioedema; LS=least squares
Note: LS means were obtained from a Poisson model accounting for overdispersion with an offset term for time on treatment and fixed effects for treatment group and HAE attack rate over a 2-month period prior to treatment
^a Obtained from a two-sided Wilcoxon rank sum test at alpha = 5%

The monthly rate of HAE attacks requiring on-demand therapy and the monthly rate of moderate or severe HAE attacks were assessed as secondary endpoints in VANGUARD. There was a 91.2% reduction with ANDEMBRY relative to placebo in the monthly rate of HAE attacks requiring on-demand therapy and a 93.6% reduction with ANDEMBRY relative to placebo in the monthly rate of moderate or severe HAE attacks.
Additional secondary endpoints assessed the proportion of patients with a ≥50%, ≥70%, ≥90%, and 100% (attack-free) reduction in monthly HAE attack rate from the first dose through the end of the 6-month treatment period compared to the 2-month period prior to treatment. The proportion of patients with a ≥50%, ≥70%, ≥90%, and 100% reduction was 95%, 92%, 74%, and 62% on ANDEMBRY versus 33%, 17%, 8%, and 0% on placebo, respectively.

16 HOW SUPPLIED/STORAGE AND HANDLING

How Supplied

ANDEMBRY (garadacimab-gxii) injection is a ready-to-use, slightly opalescent to clear, brownish-yellow to yellow solution in the following presentations, supplied in single-dose prefilled autoinjector or prefilled syringe with needle safety device.
Table 5 provides the available presentations and strengths of ANDEMBRY.

Table 5 Presentations and Strengths for ANDEMBRY

Presentation	Strength	Unit Count	NDC (Device)	NDC (Carton)
Prefilled autoinjector	200 mg/1.2 mL	1	63833-925-20	63833-925-01
Prefilled syringe with needle safety device	200 mg/1.2 mL (167 mg/mL)	1	63833-920-20	63833-920-01

Each single-dose prefilled autoinjector or prefilled syringe with needle safety device is supplied in a carton.

Storage and Handling

- Do not freeze. Do not shake.
- Keep the prefilled autoinjector and prefilled syringe with needle safety device in the original carton to protect from light.
- Store the prefilled autoinjector and prefilled syringe with needle safety device refrigerated at 36°F to 46°F (2°C - 8°C).

17 PATIENT COUNSELING INFORMATION

Advise the patient to read the FDA-approved patient labeling (Patient Information and Instructions for Use).

Administration Instructions

- Instruct patients that ANDEMBRY is for subcutaneous use and intended for self-administration or administration by caregiver.
- Instruct patients and/or caregivers on the proper preparation and subcutaneous administration techniques of ANDEMBRY [see *Dosage and Administration (2.2)* and *Instructions for Use*].
- Instruct patients and/or caregivers to administer ANDEMBRY subcutaneously into the upper arm, thigh or abdomen ensuring to stay 1 inch (2 cm) away from the navel [see *Dosage and Administration (2.2)*].
- Instruct patients or caregivers in proper disposal technique for prefilled autoinjector or prefilled syringe with needle safety device and advise them not to reuse these items.
- Instruct patients to dispose of the prefilled autoinjector or prefilled syringe with needle safety device in a sharps disposal container (closed puncture-resistant container).

Manufactured for:

CSL Behring GmbH
Emil-von-Behring-Strasse 76,
35041 Marburg, Germany
US License No. 1765

Manufactured by:

CSL Behring LLC
King of Prussia, PA 19406
US License No. 1767

Distributed by:

CSL Behring LLC
Kankakee, IL 60901

For more information, visit www.ANDEMBRY.com

For Patent information: www.cslbehring.com/products/patents

PATIENT INFORMATION ANDEMBRY® (an-DEM-bree) (garadacimab-gxii) injection, for subcutaneous use

Read this Patient Information before you start using ANDEMBRY and each time you get a refill. There may be new information. This information does not take the place of talking with your healthcare provider about your medical condition or treatment.

What is ANDEMBRY?

ANDEMBRY is a prescription medicine used to prevent attacks of Hereditary Angioedema (HAE) in children 12 years of age and older. It is not known if ANDEMBRY is safe and effective in children under 12 years of age.

Before you use ANDEMBRY, tell your healthcare provider about all of your medical conditions, including if you:

- are pregnant or planning to become pregnant. It is not known if ANDEMBRY can harm your unborn baby.
- are breastfeeding or plan to breastfeed. It is not known if ANDEMBRY passes into your breastmilk. Talk to your healthcare provider about the best way to feed your baby while using ANDEMBRY.

Tell your healthcare provider about all the medicines you take, including prescription medicines, over-the-counter medicines, vitamins and herbal supplements.

How should I use ANDEMBRY?

- See the detailed Instructions for Use that comes with this Patient Information leaflet about the right way to prepare and inject ANDEMBRY.
- Use ANDEMBRY exactly as your healthcare provider tells you to use it.
- ANDEMBRY is given as an injection under your skin (subcutaneous) by you or a caregiver.
- Your healthcare provider should show you or your caregiver how to prepare and inject your dose of ANDEMBRY before you inject yourself for the first time.
- **Do not** try to inject ANDEMBRY unless you have been trained by your healthcare provider.

What are the possible side effects of ANDEMBRY?

The most common side effects of ANDEMBRY include:

- redness, itchiness, and bruising (injection site reactions)
- stomach (abdominal) pain
- runny or stuffy nose, sneezing, watery eyes (nasopharyngitis)

Tell your healthcare provider if you have any side effect that bothers you or that does not go away. These are not all the possible side effects of ANDEMBRY. For more information, ask your healthcare provider or pharmacist.

Call your doctor for medical advice about side effects. You may report side effects to FDA at 1-800-FDA-1088.

General information about the safe and effective use of ANDEMBRY

Medicines are sometimes prescribed for purposes other than those listed in a Patient Information leaflet. Do not use ANDEMBRY for a condition for which it is not prescribed. Do not give ANDEMBRY to other people, even if they have the same symptoms that you have. It may harm them. You can ask your pharmacist or healthcare provider for information about ANDEMBRY that is written for healthcare professionals.

What are the ingredients in ANDEMBRY?

Active ingredient: garadacimab-gxii

Inactive ingredients: arginine monohydrochloride, histidine, polysorbate 80, proline, water for injection.

Manufactured at:

CSL Behring GmbH
Emil-von-Behring-Strasse 76,
35041 Marburg, Germany
US License No. 1765

Manufactured by:

CSL Behring LLC
King of Prussia, PA 19406
US License No. 1767

Distributed by:

CSL Behring LLC
Kankakee, IL 60901

For Patent information: www.cslbehring.com/products/patents

For more information, visit www.ANDEMBRY.com
or call 1-866-915-6158.

This Patient Information has been approved by the U.S. Food and Drug Administration

Revised June 2025

INSTRUCTIONS FOR USE ANDEMBRY (an-DEM-bree) (garadacimab-gxii) injection, for subcutaneous use Prefilled Autoinjector

Important:

This autoinjector works differently than other injection devices. Read the Instructions for Use carefully before using it, and each time you get a new autoinjector. There may be new information. This information does not take the place of talking to your healthcare provider about your medical condition or treatment.

Children should inject ANDEMBRY under the supervision of an adult.
Make sure you have been trained by your healthcare provider before you use this autoinjector for the first time.

Parts of the Autoinjector (see Figure A):

Continue to the next sections to prepare and perform the injection.




Figure A

Read the Following Safety Information:

- Keep the autoinjector in its original carton box until use, to protect it from light.
 - Do not** remove the clear autoinjector cap until you are ready to inject.
 - Do not** put the clear autoinjector cap back on the autoinjector after it has been removed because this could start the injection and cause injury.
 - The autoinjector contains 1 dose and is for one time use only. **Do not** try to reuse the same autoinjector.
 - Do not** use the autoinjector if the expiration date has passed.
 - The autoinjector is for subcutaneous (under the skin) injection only.
 - Do not** use the autoinjector if it has dropped, looks damaged, has cracks, or is leaking medicine. Throw away the autoinjector as described in **Step 11** and use a new one.
 - Do not** inject through clothing.
 - Do not** touch or try to remove the gray needle shield at any time.
 - Keep ANDEMBRY and all medicines out of reach of children.**
- Contact your healthcare provider if you have any questions.

How Should I Store ANDEMBRY?

- Store ANDEMBRY autoinjector in a refrigerator, between 36°F to 46°F (2°C to 8°C) in its original carton box until use, to protect it from light.
- Do not** shake.
- Do not** freeze. If the autoinjector has been frozen, **do not** use the autoinjector even if it is thawed.
- The refrigerated autoinjector may be used until the expiration date (EXP) printed on the label.
- Take the autoinjector out of the refrigerator 30 minutes before use, allowing it to reach room temperature.
- Do not** put the autoinjector back in the refrigerator after it has reached room temperature.

Supplies needed for your Autoinjector Injection (see Figure B):

Included in the carton box:

- 1 Single-dose autoinjector

Required supplies but not included in the carton box:

- 1 Alcohol pad
- 1 Cotton ball or gauze pad
- 1 FDA-cleared sharps container or puncture-resistant container for disposal (see **Step 11. Disposing of the Autoinjector**)




Figure B

Preparing for an Injection

Do not remove the clear autoinjector cap until immediately before the injection.

Step 1. Let the Autoinjector Reach Room Temperature

- Remove the autoinjector from the carton box and place it laying down-on a clean flat surface.
- Wait **30 minutes** for the medicine to reach room temperature after taking it out of the refrigerator (see **Figure C**).
- Injecting the medicine cold could be uncomfortable.
- Do not** try to speed up the warming process in any way. For example, **do not** warm it in a microwave, in hot water, or leave it in direct sunlight.




Figure C

Step 2. Check the Expiration Date

- Check the expiration date on the autoinjector label (see **Figure D**).
- Do not** use the autoinjector if the expiration date has passed.
- If the expiration date has passed, then safely dispose of the autoinjector and get a new one (see **Step 11. Disposing of the Autoinjector**).




Figure D

Step 3. Inspect the autoinjector and the medicine

- **Check** the autoinjector **for damage**.
- **Check the medicine** through the viewing window of the autoinjector (see **Figure E**).
- It is normal to see air bubbles, **do not** try to remove the air bubbles.
- The medicine should be brownish-yellow to yellow and may appear slightly opalescent to clear.
- **Do not use** the autoinjector, safely dispose and get a new one (see **Step 11. Disposing of the Autoinjector**) if:
 - The medicine is discolored or contains particles
 - The autoinjector looks damaged or has cracks
 - The autoinjector is leaking
 - The autoinjector has been dropped on a hard surface, even if it does not look damaged




Figure E

Choose and Prepare an Injection Site

Step 4. Clean your Hands

- Wash your hands well with soap and water or use hand sanitizer (see **Figure F**).
-




Figure F

Step 5. Select the Injection Site

- Inject into the **thigh or belly (abdomen)** area, but stay 1 inch (2 cm) away from the belly button (navel) (see **Figure G**).
- If somebody else (caregiver) gives you the injection, they can also use the upper arm. **Do not** try to inject into the upper arm yourself.
- Change (rotate) your injection site with each injection. **Do not inject** in the same place multiple times if the skin is damaged.
- **Do not** inject into the belly button, moles, scars or bruises, or into areas where the skin is tender, red, hard or injured.




Figure G

Step 6. Prepare the Injection Site

- Clean the injection site with an alcohol pad (see **Figure H**).
- Let your skin dry on its own.
- **Do not** touch this area again before injecting.
- **Do not** fan or blow on the skin area that you cleaned.




Figure H

Injecting the Medicine with the Autoinjector

Complete the Injection without stopping.

Read all steps first before beginning.

Do not remove the Clear Cap until you are ready to inject.

Step 7. Remove Clear Autoinjector Cap and Dispose of the Cap

- Hold the autoinjector with one hand and **pull the Clear Autoinjector Cap straight off** with the other hand.
- **Do not** twist the Clear Cap, (see **Figure I**). If you cannot remove the Clear Cap, ask a caregiver for help or contact your healthcare provider.
- The Clear Cap has a metal part inside, this is normal.
- **Do not put the Clear Cap back on** after it has been removed, because this could start the injection and cause injury.
- Dispose of the Clear Cap in an FDA-cleared sharps container.

Important:

Do not touch the gray needle shield of the autoinjector to avoid injury.

Do not put the autoinjector down after removing the Clear Cap.




Figure I

Step 8. Pinch the Skin and place the autoinjector on the injection site

Immediately after removing the clear autoinjector cap, complete the following steps without stopping:

- **Gently pinch** the area of cleaned skin around the injection site and hold the area firmly until the injection is complete (see **Figure J**).
- Place the autoinjector at a **90° angle** on the cleaned injection site (see **Figure J**).
- **Make sure you can see the viewing window.**




Figure J

Step 9. Inject Medicine (see Figure K)



You Must Read all of Step 9 before injecting.
The injection may take up to 15 seconds.

To ensure you receive a full dose you must keep the autoinjector firmly pressed against your pinched skin until:

- **The yellow plunger has stopped moving and filled the viewing window,** and
- **5 seconds has passed after the 2nd "Click."**

Press the gray needle shield down firmly against the pinched skin to start the injection and keep pressing down until all the steps below are complete.


Press down to start	Keep Pressing Down	Keep Pressing down for 5 more seconds ...1...2...3...4...5
		
Press down to start the injection and listen for a First 'Click' .	Keep pressing the autoinjector down and watch the viewing window.	Keep pressing the autoinjector down for 5 more seconds to make sure you get the full dose.
• The First "Click" means the injection has started. • The yellow plunger will start moving in the viewing window. Keep pressing the autoinjector down.	• The window will turn yellow and • You will hear a Second 'Click'. Keep pressing the autoinjector down.	Make sure the yellow plunger completely fills the window and has stopped moving before you remove the autoinjector.

Figure K

- **Do not** remove the autoinjector until the yellow plunger has stopped moving and completely filled the viewing window, and 5 seconds has passed after the Second "Click."
- **Do not** remove, tilt, or rotate the autoinjector during the injection.

Step 10. Release the Pinch and Remove the Autoinjector

- Release the pinch and remove the autoinjector at a 90° angle from the skin (see **Figure L**).
- As the autoinjector is lifted from the skin, the gray needle shield returns to the original (before use) position, and will lock into place, covering the needle.




Figure L

Important: If you think that you have not received the full dose contact your healthcare provider right away.

- If there is a little bleeding at the injection site, you can press a cotton ball or gauze over the injection site.
- **Do not** rub the injection site.
- If needed, you may cover the injection site with a small adhesive bandage.

Step 11. Disposing of the autoinjector

- **Do not** try to reuse the autoinjector.
- After injecting your dose, put the autoinjector into an FDA-cleared sharps container or closed puncture-resistant container (see **Figure M**).



Figure M

- If you do not have an FDA-cleared sharps container, you may use a household container that is:
 - Made of heavy-duty plastic
 - Can be closed with a tight-fitting, puncture-resistant lid, without sharps being able to come out
 - Upright stable during use
 - Leak-resistant
 - Properly labeled to warn of hazardous waste inside the container
- When your sharps container is almost full, you will need to follow your local guidelines for the right way to dispose of your sharps container. There may be state or local laws about how you should throw away used needles and syringes. For more information, go to the FDA's website at: <http://www.fda.gov/safesharpsdisposal>.
- **Do not** dispose of your used sharps container in your household trash unless your local guidelines permit this.
- **Do not** recycle your used sharps container.

Step 12. Keep Track of Your Treatment

- If required by your healthcare provider, record your injection in a diary to help keep track of your medicine.

Manufactured by:

CSL Behring LLC
King of Prussia, PA 19406
US License No. 1767

This Instructions for Use has been approved by the U.S. Food and Drug Administration.
Revised: June 2025